





















#Not actual size

\*Each drug is recommended for use in combination with other antiretrovirals. Please refer to the full Prescribing Information for each medication for complete indication, efficacy, safety and dosing considerations.

Drug Class	Brand (Generic Name)	Adult Daily Dosing#	Adult Regimen*	Food and liquid qualifications
Protease Inhibitors (PI)	Agenerase (amprenavir)		Eight 150-mg capsules, twice daily	May be taken with or without food, but a high fat meal decreases absorption and should be avoided.
	Crixivan (indinavir)		Two 400-mg capsules, every 8 hours	Take one hour before or two hours after meal with water; consume 1.5 liters of water a day
	Fortovase (saquinavir soft-gel)		Six 200-mg capsules, three times a day	Take with meal or within 2 hours after a meal
	Kaletra (lopinavir/ritonavir)		Three capsules (133.3-mg lopinavir/33.3-mg ritonavir), twice daily	Take with food
	Norvir (ritonavir)		Six 100-mg capsules, twice daily (start with 300-mg twice daily and increase by 100-mg twice daily every 2-3 days)	Take with food, if possible
	Viracept (nelfinavir)		Five 250-mg tablets, twice daily (or three 250-mg tablets three times a day)	Take with a meal or light snack
Nucleoside Reverse Transcriptase Inhibitor (NRTI)	Combivir (lamivudine/zidovudine)		One tablet (150-mg lamivudine, 300-mg zidovudine), twice daily	No food restrictions
	Epivir (lamivudine, 3TC)		One 150-mg tablet, twice daily	No food restrictions
	Hivid (zalcitabine, ddC)		One 0.75-mg tablet, three times a day	No food restrictions
	Retrovir (zidovudine, AZT)		One 300-mg tablet, twice daily	No food restrictions
	Trizivir (abacavir/ lamivudine/ zidovudine)		One tablet (300-mg abacavir, 150-mg lamivudine, 300-mg zidovudine), twice daily	No food restrictions
	Videx EC (didanosine, ddI)		One 400-mg capsule, daily for adults weighing 132lbs/60kg or more	Take on an empty stomach, at least 30 minutes before or 2 hours after a meal
	Zerit (stavudine, d4T)		One 40-mg capsule, twice daily for adults weighing 132lbs/60kg or more	No food restrictions
	Ziagen (abacavir sulfate)		One 300-mg tablet, twice daily	No food restrictions
Nucleotide (NRTI)	Viread (tenofovir)		One 300-mg tablet daily	Take with food
Non-nucleoside Reverse Transcriptase Inhibitor (NNRTI)	Rescriptor (delavirdine)		Two 200-mg tablets, three times a day	No food restrictions (take one hour apart from antacids)
	Sustiva (efavirenz)		Three 200-mg capsules daily	Avoid taking with high fat meal
	Viramune (nevirapine)		One 200-mg tablet daily for the first 14 days, then one 200-mg tablet, twice daily	No food restrictions