

Research article

Open Access

Demographic, risk behaviour and personal network variables associated with prevalent hepatitis C, hepatitis B, and HIV infection in injection drug users in Winnipeg, Canada

John L Wylie*^{1,2,3}, Lena Shah² and Ann M Jolly^{4,5}

Address: ¹Department of Medical Microbiology, University of Manitoba, Winnipeg, Manitoba, Canada, ²Department of Community Health Sciences, University of Manitoba, Winnipeg, Manitoba, Canada, ³Cadham Provincial Laboratory, Manitoba Health, Winnipeg, Manitoba, Canada, ⁴The Centre for Infectious Disease Prevention and Control, Population and Public Health Branch, Health Canada, Ottawa, Ontario, Canada and ⁵Department of Epidemiology and Community Medicine, University of Ottawa, Ottawa, Ontario, Canada

Email: John L Wylie* - JWylie@gov.mb.ca; Lena Shah - Lena.Shah@moh.gov.on.ca; Ann M Jolly - ann_m_jolly@phac-aspc.gc.ca

* Corresponding author

Published: 13 September 2006

Received: 05 June 2006

BMC Public Health 2006, **6**:229 doi:10.1186/1471-2458-6-229

Accepted: 13 September 2006

This article is available from: <http://www.biomedcentral.com/1471-2458/6/229>

© 2006 Wylie et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: Previous studies have used social network variables to improve our understanding of HIV transmission. Similar analytic approaches have not been undertaken for hepatitis C (HCV) or B (HBV), nor used to conduct comparative studies on these pathogens within a single setting.

Methods: A cross-sectional survey consisting of a questionnaire and blood sample was conducted on injection drug users in Winnipeg between December 2003 and September 2004. Logistic regression analyses were used to correlate respondent and personal network data with HCV, HBV and HIV prevalence.

Results: At the multivariate level, pathogen prevalence was correlated with both respondent and IDU risk network variables. Pathogen transmission was associated with several distinct types of high-risk networks formed around specific venues (shooting galleries, hotels) or within users who are linked by their drug use preferences. Smaller, isolated pockets of IDUs also appear to exist within the larger population where behavioural patterns pose a lesser risk, unless or until, a given pathogen enters those networks.

Conclusion: The findings suggest that consideration of both respondent and personal network variables can assist in understanding the transmission patterns of HCV, HBV, and HIV. It is important to assess these effects for multiple pathogens within one setting as the associations identified and the direction of those associations can differ between pathogens.

Background

The likelihood of exposure to bloodborne pathogens is a multifactorial process primarily dependent on the risk behaviours an individual practices and the likelihood that a susceptible individual will come into contact with an infected individual (thus increasing or decreasing the risk

actually associated with a risk behaviour). The interactions that bring susceptible and infected individuals into contact with each other occur within the context of social networks, the overall structure of which can also affect the rate of pathogen spread [1,2]. Chance events also play a role as a social network may contain a group of individu-

als whose behaviours favour transmission, but until such time as a bloodborne pathogen enters that network, they face no risk of acquiring that pathogen.

Individual risk behaviours for injection drug users (IDU) include those directly associated with transmission, such as the use of syringes previously used by another IDU or those behaviours which can act as markers of the above types of behaviours. Examples of risk markers that are positively associated with disease prevalence include drug scene roles, such as dealing drugs or injecting others as a service (street or hit doctors) [3-6]. Other behaviours, such as obtaining clean needles from questionable sources such as drug dealers, shooting gallery owners, or on the street, can also act as a marker of an increased probability of using a contaminated needle, as doctoring of used needles to make them appear new has been reported [7].

Social network analysis is a technique which measures and analyzes the interactions that occur between people and its application can enhance an otherwise overly simplistic interpretation of individual risk behaviours. In Brooklyn, injecting with a used syringe was associated with HIV only if syringes were being shared with someone who was 10 or more years older than the interviewed subject, or was a daily injector [8]. By incorporating network variables indicative of a higher probability of coming into contact with an infected individual, a better indication is obtained of who is at risk of infection and what actually constitutes a high-risk behaviour.

Subsequent research on IDU and their network characteristics have identified other network variables associated with transmission risk. High-risk injection practices have been linked to network characteristics such as the number of network members [9,10]; the presence of family members or spouses within the network [4,11]; higher network density [10]; the setting where injection takes place [10]; turnover of network members [12]; and the pooling of financial resources within networks for the purpose of obtaining drugs [9]. Racial/ethnic differences in HIV prevalence have also been at least partially explained by taking into account the differing network characteristics of different ethnic groups [13].

Some individual behaviours or characteristics may also be proxy markers of network behaviours. The type of drug an IDU chooses to inject can be measured as a characteristic of the individual injector and itself can influence risk, as some drugs, like cocaine, are prepared at room temperature and hence are more conducive to pathogen survival [14]. In addition to these more risky drug-specific practices, IDU may form networks based on drug type, which mark the broader social network within which an individ-

ual is a member [15,16]. Therefore, network members are more likely to come into contact with whichever pathogen(s) happen to be circulating within that network. Similarly, moving to a new city within the past year can also be an indicator of higher risk as individuals create a social bond through the sharing of drug equipment to try and establish themselves in new networks [12,17].

We have recently described networks centered on downtown hotels in the core area of Winnipeg (manuscript in press), which demonstrate that the setting where an individual injects drugs can also act as a proxy marker of a social network in which some of the members may not know each other, but are nonetheless linked by a common venue. Within these venues accidental or intentional use of contaminated syringes is more likely. Injection at a shooting gallery is another example of a geographic place-based network that has also frequently been linked to increased risk of infection by a bloodborne pathogen [6,18,19].

The majority of the network research described above has focused on HIV and relatively little on hepatitis B and C (HBV and HCV, respectively). Comparative studies of network variables associated with the transmission of HIV, HCV and HBV within the same setting have also not been conducted. Our goal was to determine whether respondent and network variables (including some proxy markers of networks as described above) are associated with the prevalence of HCV, HBV, and HIV in our setting and to what extent similar or dissimilar patterns emerge for the different pathogens.

Methods

Study setting

A cross-sectional survey of IDU in Winnipeg, Manitoba, Canada (pop. 675,000) from December, 2003 to September, 2004 was conducted. Potential study recruits were invited to participate through advertisements posted at local community health centres, and meeting places such as laundromats, which had been suggested by key informants within the target population. Word-of-mouth advertising also occurred in the community as the study progressed.

Eligibility criteria were self-reported use of illicit injection drugs in the 6 month period prior to interview and age 15 years or more. Potential participants made telephone contact with the study nurse, who administered all the surveys and collected a blood specimen. An honorarium of \$40 Cdn was provided to all study participants, regardless of whether they completed all parts of the study. Participants provided written or oral consent. The study design was approved by the Health Research Ethics Board of the Uni-

versity of Manitoba and the Winnipeg Regional Health Authority Research Review Committee.

Survey instrument

The questionnaire was divided into three sections. Section 1 consisted of questions based on the respondent's own demographic or behavioural characteristics (hereafter referred to as respondent variables).

Section 2 elicited information on the study participant's egocentric network. Study participants were asked to think back over the last 30 days about the people with whom they had had more than casual contact. Prompts included friends, relatives, or other individuals to whom they feel close, and people with whom they had used drugs, had sex, resided, or hung out. Using initials or other anonymous identifiers, participants were asked to list a maximum of 20 members of their egocentric network (referred to as egocentric variables).

Section 3 elicited information on the respondent's IDU risk network. In this section, a series of detailed questions were asked about each of the individuals within the egocentric network that had been identified by the respondent as IDU (to a maximum of 5 IDU). If more than 5 IDU were listed in the egocentric network, only the first 5 IDU that were listed in the egocentric network were chosen. Respondents then were asked questions about each member of his or her IDU network (e.g. have you ever used a syringe after [person] used it first?). Respondents could then provide different responses for each of the IDU within their risk network (these variables are referred to IDU risk network variables).

Measures

Respondent sociodemographic variables

Demographic variables included age, age of initiation of injection drug use (analyzed as number of years a respondent had injected drugs; referred to as "years of ID use"), gender, level of education completed, ethnicity and whether respondents had moved to Winnipeg in the past year.

Preliminary examination of the data indicated age and years of ID use were correlated; years of ID use only was chosen for use in this analysis. The relationship between years of ID use and serostatus was nonlinear (examined with Lowess smoothed plots [20]). In general, during the first years of drug use, there was a rapid increase in serostatus, followed by a more gradual increase until approximately year 20 after which it leveled off (data not shown). These breakpoints approximated the quartile distribution of years of ID use, therefore this variable was recoded to quartiles.

Due to small sample size, transgender persons ($n = 5$) and those reporting an ethnicity other than Caucasian or Aboriginal ($n = 10$) were excluded from analysis. Educational level was coded to two categories: individuals who dropped out of school and those who graduated grade 12 or were currently in school. A recent move to Winnipeg was categorized as those respondents who had moved to Winnipeg in the past year.

Respondent drug-related behaviours

Respondents were asked questions about their drug-related behaviours, which generally covered the 6 month period prior to interview, but exceptions are noted. Three variables represented use of the three most common drugs: cocaine, a combination of talwin and ritalin (talwin/ritalin) and morphine. A fourth variable for heroin injection (a relatively rare drug in Winnipeg) was included for means of comparison with other studies and because anecdotal evidence suggested that heroin users may be somewhat isolated from other types of users in Winnipeg forming a distinct network with distinct properties. Frequency of drug use was coded as daily vs. non-daily use.

Two binary variables contained data on injection at hotels or shooting galleries. Data on whether respondents had used anyone else's used syringes (ever and during the last 6 months); other related drug-preparation equipment (last 6 months; prompts included cooker, rinse water, or cotton); engaged in drug transfer behaviours (prompts included backloading, frontloading or piggybacking) from another user's syringe to their own (last 6 months) were evaluated. Univariate analysis showed that only ever-use of previously used syringes was significant and is the only one of the above variables presented in this paper (e.g. the univariate OR relating HBV serostatus and use of other's equipment was 0.98 [0.62, 1.55] while the univariate OR for HBV serostatus and engaging in drug transfer behaviours was 0.94 [0.53, 1.67]). Injecting someone as a service was defined as someone who reported that he or she had ever received any money, drugs, or other goods in exchange for injecting someone with drugs.

Obtaining needles from questionable sources such drug dealers, shooting gallery owners, or syringes found on the street was evaluated separately from those who obtained needles from friends or family members, both in the last six months. Both were constructed as binary variables.

While sexual behaviours are linked to HBV and HIV transmission, they are not considered an important route of transmission for HCV, but still indicate certain roles or networks which may be relevant and predictive of risk. We included six binary variables denoting opposite-sex or same-sex sex partnerships in the last six months of the fol-

lowing type; regular (someone with whom the participant is emotionally involved with), casual (someone with whom the participant has had sex with only a few times and has no emotional involvement with) or client (someone who has given the participant money or goods in exchange for sex).

Egocentric network variables

We created two continuous variables from the total number of people on the network list whom the participant identified as IDU and the total number of people on the list who were identified as both IDU and family members.

IDU risk network variables

A series of variables were based on the interactions which participants had with individual IDU within their egocentric network (maximum five IDU as noted above). For example, a participant could indicate that he or she saw daily anywhere from none to five of the IDU on their IDU risk network list. To avoid small cell sizes, data were collapsed to either three (0, 1, or 2–5 IDU) or four (0, 1, 2, or 3–5 IDU) dummy variable categories. Twelve interactions with other IDU were considered, including how many IDU risk members 1) the participant sees on a daily basis, 2) inject daily, 3) have been IDU for more than 5 years, 4) have injected at a hotel, 5) have injected at a shooting gallery, 6) the participant has pooled resources with to obtain drugs, 7) have used a syringe before the participant has used it, 8) have used other drug preparation equipment before the participant has used it, 9) have initiated the respondent to drug use, 10) have shown the respondent how to inject drugs, 11) have injected the respondent with drugs, 12) use talwin/ritalin, 13) cocaine, or 14) morphine (heroin was not included here as only 18 respondents knew any heroin users). Variables 7 and 8 are similar to respondent variables regarding the sharing of syringes and equipment (across all partners), however, we felt asking the questions within a network context may result in better recall of these events. Similarly, given the importance of drug type and infection risk, we included variables 12–14.

Diagnostic testing

Venous blood specimens were collected for HCV, HBV and HIV serological testing at Cadham Provincial Laboratory (Winnipeg, MB). Specimens were screened for HCV and HIV with AxSYM HCV (Abbott, Mississauga, ON) and AxSYM HIV1/2 gO (Abbott, Mississauga, ON), respectively. Presumptive positives were confirmed for HCV with Chiron HCV 3.0 RIBA (Ortho-Clinical Diagnostics, Markham, ON). Presumptive HIV positive specimens were confirmed by western blot (BioRad, Montreal, QC). HBV cases were considered to be those specimens positive

for antibodies against HBV core protein (IMX HBV core IgG, Abbott, Mississauga, ON).

Statistical analysis

Logistic regression was used to analyze the three binary dependent variables (HCV, HBV, and HIV serostatus). The multicollinearity of independent variables was first assessed with a correlation matrix [21]. As noted above, age and years of ID use were correlated and years of ID use was chosen for analysis. The only other two variables that were clearly correlated were respondent's use of talwin/ritalin and use of talwin/ritalin by IDU risk network members. Given that we were interested in comparing variables from an individual *vs.* network perspective, we did not want to choose one of these variables over the other, nor combine them into one variable. Therefore, for each pathogen, we constructed two models where either one or the other of the above variables was used. In all cases, the inclusion of one of these variables over the other had no major effect on the other independent variables that remained in the final models or on the final odds ratios (data not shown). We present only the models using IDU risk network use of talwin/ritalin as it illustrated a trend toward increasing prevalence for some pathogens as the number of talwin/ritalin users within the risk network increased. All remaining variables that were entered into multivariable analysis were also checked for multicollinearity using NCSS (Kaysville, UT). Variance inflation factor and tolerance statistics were within acceptable limits for all variables.

Univariate analyses were first completed and all variables with *p* values of 0.2 or less were considered for inclusion in multivariable analyses. Logistic regression model building procedures followed those described by Hosmer and Lemeshow [20]. Multivariable analysis was first conducted by entering respondent variables into a regression model. The effect of removing individual variables was assessed through the likelihood ratio test. After final selection of respondent variables, egocentric and IDU risk network variables were entered and the effect of removing each respondent and network variable was again tested with the likelihood ratio test. After creating preliminary main effects models, interactions were assessed between the respondent variable, injection with a used syringe, and all other variables in the model. These interactions were assessed to determine whether any of the variables in the model may affect the risk associated with syringe sharing. All statistical analysis was done using STATA version 8 (Stata corporation, College Station, TX). Diagnostics for the final models were performed using the Hosmer-Lemeshow goodness of fit statistic and the area under the receiver operating characteristic curve using the *lfit* and *lroc* commands in STATA [20].

Results

A total of 435 IDU were enrolled in the study. Some IDU were unable or unwilling to provide a blood specimen. Additionally, five individuals who self-identified as transgender and 10 individuals who self-identified as belonging to ethnic groups other than Caucasian or Aboriginal, were excluded from the analysis. For the remaining 369 respondents, summary statistics for the egocentric network are as follows: mean number of members, 8.8; median, 8; range, 1–20, lower quartile, 5; upper quartile, 11; SD, 4.6. Given that the IDU risk network was truncated at 5 IDU, we also report these summary statistics for the total number of people within the egocentric network identified as IDU: mean number of IDU in the egocentric network, 3.5; median, 3; range, 0–20; lower quartile, 1; upper quartile, 5. In total, 17% of respondents (66/369) reported more than 5 IDU in their egocentric network. The lack of data on these individuals is discussed further in the limitations section of this paper. Of the 369 respondents considered for analysis, several indeterminate results were identified for each pathogen therefore, the final sample sizes for HCV, HBV, and HIV were 365, 364, and 360, respectively.

Correlation between HCV, HBV, and HIV

To assist in data interpretation the correlation between HCV, HBV, and HIV were determined. Pairwise correlation values for HCV/HBV serostatus was 0.001; HCV/HIV, -0.011; and HBV/HIV, -0.011. No apparent correlation between the three pathogens was evident.

Univariate analyses

The univariate analyses for these 3 pathogens are shown in tables 1, 2, 3. In univariate analyses, the subset of variables associated with serostatus and the direction of that association was similar for the three pathogens. Respondent variables positively associated with serostatus for all three pathogens were: years of ID use, aboriginal ethnicity, injection at a shooting gallery, ever reporting the use of someone's else's used syringe; injecting someone else as a service; and having opposite-sex client partners.

Thirteen variables were associated with both HCV and HBV, but not HIV. Positive associations were found for two respondent variables (having injected at a hotel; injecting talwin/ritalin), two egocentric network variables (total number of IDU; total number of family IDU), and five IDU risk network variables (the number of IDU who inject daily, the number of IDU who have been injecting for more than 5 years, the number of IDU who have injected talwin/ritalin, the number of IDU who have injected at a hotel, and the number of IDU who have used a syringe before the respondent used it). Negative associations were found for three respondent variables (moved to Winnipeg in the previous year; injection of heroin and

daily injection) and one IDU risk network variable (number of IDU who initiated the respondent to injection drug use).

Two respondent variables were associated with HCV and HIV, but not HBV. A negative association was found for obtaining clean syringes from friends. Reporting same-sex regular sex partners was negatively associated with HCV and positively associated with HIV.

Ten variables were associated with only one of the three pathogens. Positive associations were found between HCV and one respondent variable (obtains clean syringes from questionable sources); between HBV and four IDU risk network variables (the number of IDU in the risk network who have injected at a shooting gallery; the number of IDU with whom the respondent has pooled resources; the number of IDU who have injected the respondent and the number of IDU who have injected cocaine); between HIV and two respondent variables (injection of cocaine and same-sex client sex partners). Negative associations were found between HBV and reporting same-sex casual sex partners and opposite-sex casual partners and between HCV and the number of IDU in the risk network who have shown the respondent how to inject drugs.

Multivariate analyses

In multivariate analysis, prior to assessing interactions, HCV was positively associated with respondent's age of initiation to injection drug use, reported injection with a syringe used by another IDU, injection at a shooting gallery, injecting someone as a service, and the number of IDU in the risk network who inject talwin/ritalin (table 4). Negative associations for this pathogen were found for the respondent obtaining syringes from friends and the number of IDU in the risk network who had shown the respondent how to inject drugs.

For the model described above, interactions were assessed between the respondent variable, injection with a used syringe, and the remaining variables in the model. An interaction was noted between injection with used syringes and the number of IDU network members who inject talwin/ritalin and the number of IDU network members who had shown the respondent how to inject drugs (table 5). For respondents who did not associate with other talwin/ritalin users, injecting with syringes previously used by other IDUs was positively associated with HCV prevalence. This correlation was absent among those IDU linked to talwin and ritalin, such that their HCV prevalence was high regardless of whether or not they reported injecting with used syringes. Similarly, for those IDU who were in a risk network where someone had shown them how to inject drugs, HCV prevalence was relatively low and not associated with injection with a used syringe.

Table 1: Odds ratios and 95% confidence intervals for univariate analysis of HCV serostatus.

Variable name	HCV negative (n = 167) (%)	HCV positive (n = 198) (%)	OR (95% CI)	p value
Respondent variables				
Years of injection drug use (quartiles)				
0–6	81 (48.5)	20 (10.1)	1.0	
7–13	46 (27.5)	51 (25.8)	4.5 (2.4–8.4)	<0.001
14–21	19 (11.4)	61 (30.8)	13.0 (6.4–26.5)	<0.001
22–39	21 (12.6)	66 (33.3)	12.7 (6.4–25.5)	<0.001
Gender				
Male	96 (57.5)	113 (57.1)	1.0	
Female	71 (42.5)	85 (42.9)	1.0 (0.7–1.5)	0.937
Education level				
< grade 12	116 (69.5)	142 (71.7)	1.0	
> grade 12	51 (30.5)	56 (28.3)	0.9 (0.6–1.4)	0.637
Ethnicity				
Caucasian	71 (42.5)	57 (28.8)	1.0	
Aboriginal	96 (57.5)	141 (71.2)	1.8 (1.2–2.8)	0.006
Moved to Winnipeg in past year				
No	112 (67.1)	153 (77.3)	1.0	
Yes	55 (32.9)	45 (22.7)	0.6 (0.4–1.0)	0.030
Deals drugs				
No	85 (50.9)	107 (54.0)	1.0	
Yes	82 (49.1)	91 (46.0)	0.9 (0.6–1.3)	0.549
Cocaine use				
No	62 (37.1)	69 (34.9)	1.0	
Yes	105 (62.9)	129 (65.1)	1.1 (0.7–1.7)	0.651
Talwin/ritalin use				
No	134 (80.2)	113 (57.1)	1.0	
Yes	33 (19.8)	85 (42.9)	3.1 (1.9–4.9)	<0.001
Morphine use				
No	110 (65.9)	121 (61.1)	1.0	
Yes	57 (34.1)	77 (38.9)	1.2 (0.8–1.9)	0.347
Heroin use				
No	145 (86.8)	184 (92.9)	1.0	
Yes	22 (13.2)	14 (7.1)	0.5 (0.2–1.0)	0.052
Injects daily				
No	20 (12.0)	43 (21.7)	1.0	
Yes	147 (88.0)	155 (78.3)	0.5 (0.3–0.9)	0.013
Injected at a hotel				
No	109 (65.3)	107 (54.0)	1.0	
Yes	58 (34.7)	91 (46.0)	1.6 (1.0–2.4)	0.029
Injected at a shooting gallery				
No	153 (91.6)	156 (78.8)	1.0	
Yes	14 (8.4)	42 (21.2)	2.9 (1.5–5.6)	<0.001
Ever used someone else's syringe				
No	90 (53.9)	49 (24.8)	1.0	
Yes	77 (46.1)	149 (75.3)	3.6 (2.3–5.5)	<0.001
Injected someone else as a service				

Table 1: Odds ratios and 95% confidence intervals for univariate analysis of HCV serostatus. (Continued)

No	144 (86.2)	125 (63.1)	1.0	
Yes	23 (13.8)	73 (36.9)	3.7 (2.2–6.2)	<0.001
Obtains syringes from friends				
No	50 (29.9)	73 (36.9)	1.0	
Yes	117 (70.1)	125 (63.1)	0.7 (0.5–1.1)	0.162
Obtains syringes from questionable sources				
No	135 (80.8)	144 (72.7)	1.0	
Yes	32 (19.2)	54 (27.3)	1.6 (1.0–2.6)	0.067
Opposite-sex regular partners				
No	51 (30.5)	65 (31.8)	1.0	
Yes	116 (69.5)	163 (68.2)	0.9 (0.6–1.4)	0.640
Opposite-sex casual partners				
No	101 (60.5)	129 (65.2)	1.0	
Yes	66 (39.5)	69 (34.9)	0.8 (0.5–1.3)	0.357
Opposite-sex client partners				
No	151 (90.4)	163 (82.3)	1.0	
Yes	16 (9.6)	35 (17.7)	2.0 (1.1–3.8)	0.024
Regular same-sex partners				
No	160 (95.8)	195 (98.5)	1.0	
Yes	7 (4.2)	3 (1.5)	0.4 (0.1–1.4)	0.116
Casual same-sex partners				
No	160 (95.8)	191 (96.5)	1.0	
Yes	7 (4.2)	7 (3.5)	0.8 (0.3–2.4)	0.745
Client same-sex partners				
No	163 (97.6)	194 (98.0)	1.0	
Yes	4 (2.4)	4 (2.0)	0.8 (0.2–3.4)	0.808
Egocentric network variables (maximum number of network members = 20)				
Total number of IDU in network				
Median (IQR)	3 (1,5)	3 (1,5)	1.0 (1.0–1.1)	0.194
Total number of IDU family members in network				
Median (IQR)	0 (0,0)	0 (0,1)	1.29 (1.1–1.6)	0.10
Spouse IDU in network				
No	109 (65.3)	127 (64.1)	1.0	
Yes	58 (34.7)	71 (35.9)	1.1 (0.7–1.6)	0.822
IDU risk network variables (maximum number of IDU = 5)				
Number of IDU respondent sees daily				
None	54 (32.3)	66 (33.3)	1.0	
1	55 (32.9)	68 (34.3)	1.0 (0.6–1.7)	0.964
2	32 (19.2)	30 (15.2)	0.8 (0.4–1.4)	0.398
3–5	26 (15.6)	34 (17.2)	1.1 (0.6–2.0)	0.832
Number of IDU in network who inject daily				
None	104 (62.3)	105 (53.0)	1.0	
1	32 (19.2)	36 (18.2)	1.1 (0.6–1.9)	0.699
2	19 (11.4)	31 (15.7)	1.6 (0.9–3.0)	0.137
3–5	12 (7.2)	26 (13.1)	2.1 (1.0–4.5)	0.042
Number of IDU in network who have been IDU for >5 years				
0	101 (60.5)	74 (37.4)	1.0	
1	28 (16.8)	40 (20.2)	1.9 (1.1–3.4)	0.021

Table 1: Odds ratios and 95% confidence intervals for univariate analysis of HCV serostatus. (Continued)

2	18 (10.8)	39 (19.7)	3.0 (1.6–5.6)	0.001
3–5	20 (12.0)	45 (22.7)	3.1 (1.7–5.6)	<0.001
Number of IDU in network who have injected at a hotel				
0	117 (70.1)	112 (56.6)	1.0	
1	29 (17.4)	39 (19.7)	1.4 (0.8–2.4)	0.222
2–5	21 (12.6)	47 (23.7)	2.3 (1.3–4.2)	0.004
Number of IDU in network who have injected at a shooting gallery				
0	134 (80.2)	143 (72.2)	1.0	
1	18 (10.8)	27 (13.6)	1.4 (0.7–2.7)	0.298
2–5	15 (9.0)	28 (14.1)	1.7 (0.9–3.4)	0.102
Number of IDU in network with whom respondent has pooled money				
0	51 (30.5)	56 (28.3)	1.0	
1	42 (25.2)	54 (27.3)	1.2 (0.7–2.0)	0.576
2–3	43 (25.8)	55 (27.8)	1.2 (0.7–2.0)	0.587
4–5	31 (18.6)	33 (16.7)	1.0 (0.5–1.8)	0.922
Number of IDU in network who have used a syringe before the respondent used it				
0	133 (79.6)	146 (73.7)	1.0	
1	27 (16.2)	32 (16.2)	1.1 (0.6–1.9)	0.790
2–5	7 (4.2)	20 (10.1)	2.6 (1.1–6.4)	0.036
Number of IDU in network who have used other drug paraphernalia before respondent has used it.				
0	105 (62.9)	119 (60.1)	1.0	
1	29 (17.4)	31 (16.7)	0.9 (0.5–1.7)	0.841
2–5	33 (19.8)	48 (24.2)	1.3 (0.8–2.1)	0.342
Number of IDU in network who have initiated respondent to injection drug use				
0	110 (65.9)	157 (79.3)	1.0	
1	48 (28.8)	35 (17.7)	0.5 (0.3–0.8)	0.008
2–5	9 (5.4)	6 (3.0)	0.5 (0.2–1.4)	0.160
Number of IDU in network who have shown respondent how to inject drugs				
0	104 (62.3)	147 (74.2)	1.0	
1	43 (25.8)	39 (19.7)	0.6 (0.4–1.1)	0.083
2–5	20 (12.0)	12 (6.1)	0.4 (0.2–0.9)	0.027
Number of IDU in network who have injected respondent with drugs				
0	89 (53.3)	88 (44.4)	1.0	
1	44 (26.4)	54 (27.3)	1.2 (0.8–2.0)	0.392
2–5	34 (20.4)	56 (28.3)	1.7 (1.0–2.8)	0.054
Number of IDU in network who have used talwin/ritalin				
0	139 (83.2)	119 (60.1)	1.0	
1–3	17 (10.1)	42 (21.1)	2.9 (1.6–5.3)	<0.001
4–5	11 (6.6)	37 (18.7)	3.9 (1.9–8.0)	<0.001
Number of IDU in network who have used cocaine				
0	113 (67.7)	120 (60.6)	1.0	
1	23 (13.8)	36 (18.2)	1.5 (0.8–2.6)	0.192
2–3	17 (10.2)	30 (15.2)	1.7 (0.9–3.2)	0.125
4–5	14 (8.4)	12 (6.1)	0.8 (0.4–1.8)	0.605
Number of IDU in network who have used morphine				
0	129 (77.3)	149 (75.3)	1.0	
1	22 (13.2)	31 (15.7)	1.2 (0.7–2.2)	0.513
2–5	16 (9.6)	18 (9.1)	1.0 (0.5–2.0)	0.942

Table 2: Odds ratios and 95% confidence intervals for univariate analysis of HBV serostatus.

Variable name	HBV negative (n = 253) (%)	HBV positive (n = 111) (%)	OR (95% CI)	p value
Respondent variables				
Years of injection drug use (quartiles)				
0–6	97 (38.3)	5 (4.5)	1.0	
7–13	74 (29.3)	23 (20.7)	6.0 (2.2–16.6)	<0.001
14–21	42 (16.6)	36 (32.4)	16.6 (6.1–45.3)	<0.001
22–39	40 (15.8)	47 (42.3)	22.8 (8.4–61.5)	<0.001
Gender				
Male	150 (59.3)	60 (54.0)	1.0	
Female	103 (40.7)	51 (46.0)	1.2 (0.8–1.9)	0.353
Education level				
< grade 12	178 (70.4)	79 (71.2)	1.0	
> grade 12	75 (29.6)	32 (28.8)	1.0 (0.6–1.3)	0.485
Ethnicity				
Caucasian	99 (39.1)	29 (26.1)	1.0	
Aboriginal	154 (60.9)	82 (73.9)	1.8 (1.1–3.0)	0.015
Mobility				
In Winnipeg > 1 year	168 (66.4)	97 (87.4)	1.0	
Moved to Winnipeg in past year	85 (33.6)	14 (12.6)	0.3 (0.2–0.5)	<0.001
Deals drugs				
No	129 (51.0)	61 (55.0)	1.0	
Yes	124 (49.0)	50 (45.1)	0.9 (0.5–1.3)	0.485
Cocaine use				
No	85 (33.6)	44 (39.6)	1.0	
Yes	168 (66.4)	67 (60.4)	0.8 (0.5–1.2)	0.27
Talwin/ritalin use				
No	190 (75.1)	55 (49.6)	1.0	
Yes	63 (24.9)	56 (50.5)	3.1 (1.9–4.9)	<0.001
Morphine use				
No	165 (64.8)	68 (61.3)	1.0	
Yes	89 (35.2)	43 (38.7)	1.2 (0.7–1.8)	0.516
Heroin use				
No	222 (87.8)	106 (95.5)	1.0	
Yes	31 (12.3)	5 (4.5)	0.3 (0.1–0.9)	0.015
Injects daily				
No	34 (13.4)	26 (23.4)	1.0	
Yes	219 (86.6)	85 (76.6)	0.5 (0.3–0.9)	0.021
Injected at a hotel				
No	160 (63.2)	53 (47.8)	1.0	
Yes	93 (36.8)	58 (52.2)	1.9 (1.2–3.0)	0.006
injected at a shooting gallery				
No	224 (88.5)	85 (62.2)	1.0	
Yes	29 (11.5)	26 (23.4)	2.4 (1.3–4.2)	0.004
Ever used someone else's syringe				
No	114 (45.1)	25 (22.5)	1.0	
Yes	139 (54.9)	86 (77.5)	2.8 (1.7–4.7)	<0.001
Injected someone else as a service				

Table 2: Odds ratios and 95% confidence intervals for univariate analysis of HBV serostatus. (Continued)

No	204 (80.6)	66 (59.5)	1.0	
Yes	49 (19.4)	45 (40.6)	2.8 (1.7–4.6)	<0.001
Obtains syringes from friends				
No	81 (32.0)	41 (16.9)	1.0	
Yes	172 (68.0)	70 (63.1)	0.8 (0.5–1.3)	0.362
Obtains syringes from questionable sources				
No	197 (77.9)	81 (73.0)	1.0	
Yes	56 (22.1)	30 (27.0)	1.3 (0.8–2.2)	0.316
Opposite-sex regular partners				
No	81 (32.0)	36 (32.4)	1.0	
Yes	172 (68.0)	75 (67.6)	1.0 (0.6–1.6)	0.938
Opposite-sex casual partners				
No	149 (58.9)	80 (72.1)	1.0	
Yes	104 (41.1)	31 (27.9)	0.6 (0.3–0.9)	0.015
Opposite-sex client partners				
No	223 (88.1)	90 (81.1)	1.0	
Yes	30 (11.9)	21 (18.9)	1.7 (0.9–3.2)	0.081
Regular same-sex partners				
No	245 (96.8)	109 (98.2)	1.0	
Yes	8 (3.2)	2 (1.8)	0.6 (0.1–2.7)	0.448
Casual same-sex partners				
No	240 (94.9)	109 (98.2)	1.0	
Yes	13 (5.1)	2 (1.8)	0.3 (0.1–1.5)	0.113
Client same-sex partners				
No	247 (97.6)	109 (98.2)	1.0	
Yes	6 (2.4)	2 (1.8)	0.8 (0.2–3.8)	0.728
Egocentric network variables (maximum number of network members = 20)				
Total number of IDU in network				
Median (IQR)	3 (1,4)	3 (2,5)	1.1 (1.0–1.2)	0.001
Total number of IDU family members in network				
Median (IQR)	0 (0,1)	0 (0,1)	1.3 (1.0–1.5)	0.019
Spouse IDU in network				
No	169 (66.8)	69 (62.2)	1.0	
Yes	84 (33.2)	42 (37.8)	1.2 (0.8–1.9)	0.394
IDU risk network variables (maximum number of IDU = 5)				
Number of IDU respondent sees daily				
None	83 (32.8)	38 (34.2)	1.0	
1	84 (33.2)	38 (34.2)	1.0 (0.6–1.7)	0.965
2	48 (19.0)	13 (11.7)	0.6 (0.3–1.2)	0.155
3–5	38 (15.0)	22 (19.8)	1.3 (0.7–2.4)	0.479
Number of IDU in network who inject daily				
None	160 (63.2)	50 (45.1)	1.0	
1	42 (16.6)	25 (22.5)	1.9 (1.1–3.4)	0.032
2	31 (12.3)	17 (15.3)	1.8 (0.9–3.4)	0.101
3–5	20 (7.9)	19 (17.1)	3.0 (1.5–6.1)	0.002
Number of IDU in network who have been IDU for >5 years				
0	142 (56.1)	34 (30.6)	1.0	
1	50 (19.8)	18 (16.2)	1.5 (0.8–2.9)	0.223

Table 2: Odds ratios and 95% confidence intervals for univariate analysis of HBV serostatus. (Continued)

2	32 (12.7)	24 (21.6)	3.1 (1.6–6.0)	0.001
3–5	29 (11.5)	35 (31.5)	5.0 (2.7–9.4)	<0.001
Number of IDU in network who have injected at a hotel				
0	175 (69.2)	54 (48.7)	1.0	
1	48 (19.0)	20 (18.0)	1.4 (0.7–2.5)	0.330
2–5	30 (11.9)	37 (33.3)	4.0 (2.3–7.1)	<0.001
Number of IDU in network who have injected at a shooting gallery				
0	204 (80.6)	74 (66.7)	1.0	
1	30 (11.9)	15 (13.5)	1.4 (0.7–2.7)	0.351
2–5	19 (7.5)	22 (19.8)	3.2 (1.6–6.2)	0.001
Number of IDU in network with whom respondent has pooled money				
0	84 (33.2)	23 (20.7)	1.0	
1	62 (24.5)	33 (29.7)	1.9 (1.0–3.6)	0.037
2–3	64 (25.3)	34 (30.6)	1.9 (1.0–3.6)	0.036
4–5	43 (17.0)	21 (18.9)	1.8 (0.9–3.6)	0.103
Number of IDU in network who have used a syringe before the respondent used it				
0	191 (75.5)	87 (78.4)	1.0	
1	47 (18.6)	12 (10.8)	0.6 (0.3–1.1)	0.097
2–5	15 (5.9)	12 (10.8)	1.8 (0.8–3.9)	0.168
Number of IDU in network who have used other drug paraphernalia before respondent has used it.				
0	155 (61.3)	67 (60.4)	1.0	
1	46 (18.2)	13 (11.8)	0.7 (0.3–1.3)	0.220
2–5	52 (20.6)	31 (27.9)	1.4 (0.8–2.3)	0.234
Number of IDU in network who have initiated respondent to injection drug use				
0	177 (70.0)	88 (79.2)	1.0	
1	63 (24.9)	21 (18.9)	0.7 (0.4–1.2)	0.159
2–5	13 (5.1)	2 (1.8)	0.3 (0.1–1.4)	0.128
Number of IDU in network who have shown respondent how to inject drugs				
0	172 (68.0)	79 (71.2)	1.0	
1	58 (22.9)	23 (20.7)	0.9 (0.5–1.5)	0.602
2–5	23 (9.1)	9 (8.1)	0.8 (0.4–1.9)	0.700
Number of IDU in network who have injected respondent with drugs				
0	133 (52.6)	43 (38.7)	1.0	
1	65 (25.7)	33 (29.7)	1.6 (0.9–2.7)	0.103
2–5	55 (21.7)	35 (31.5)	2.0 (1.1–3.4)	0.015
Number of IDU in network who have used talwin/ritalin				
0	197 (77.9)	59 (53.2)	1.0	
1–3	31 (12.3)	29 (26.1)	3.1 (1.7–5.6)	<0.001
4–5	25 (9.9)	48 (13.2)	3.1 (1.6–5.8)	0.001
Number of IDU in network who have used cocaine				
0	168 (66.4)	64 (57.7)	1.0	
1	39 (15.4)	20 (18.0)	1.3 (0.7–2.5)	0.340
2–3	26 (10.3)	21 (18.9)	2.1 (1.1–4.0)	0.022
4–5	20 (7.9)	26 (7.1)	0.8 (0.3–2.0)	0.625
Number of IDU in network who have used morphine				
0	198 (78.3)	80 (72.1)	1.0	
1	34 (13.4)	18 (16.2)	1.3 (0.7–2.5)	0.399
2–5	21 (8.3)	13 (11.7)	1.5 (0.7–3.2)	0.258

Table 3: Odds ratios and 95% confidence intervals for univariate analysis of HIV serostatus.

Variable name	HIV negative (n = 334) (%)	HIV positive (n = 26) (%)	OR (95% CI)	p value
Respondent variables				
Years of injection drug use (quartiles)				
0–6	100 (29.9)	2 (7.7)	1.0	
7–13	83 (24.9)	11 (42.3)	6.6 (1.4–30.7)	0.016
14–21	71 (21.3)	7 (26.9)	4.9 (0.9–24.4)	0.051
22–39	80 (24.0)	6 (23.1)	3.75 (0.7–19.0)	0.111
Gender				
Male	193 (57.8)	13 (50.0)	1.0	
Female	141 (42.2)	13 (50.0)	1.4 (0.6–3.0)	0.442
Education level				
< grade 12	234 (70.1)	19 (73.1)	1.0	
> grade 12	100 (29.9)	7 (26.9)	0.9 (0.4–2.1)	0.744
Ethnicity				
Caucasian	125 (37.4)	3 (11.5)	1.0	
Aboriginal	209 (62.6)	23 (88.5)	4.6 (1.4–15.6)	0.004
Mobility				
In Winnipeg > 1 year	244 (73.1)	18 (69.2)	1.0	
Moved to Winnipeg in past year	90 (27.0)	8 (30.8)	1.2 (0.5–2.9)	0.677
Deals drugs				
No	175 (52.4)	15 (57.7)	1.0	
Yes	159 (47.6)	11 (42.3)	0.8 (0.4–1.8)	0.601
Cocaine use				
No	122 (36.5)	6 (23.1)	1.0	
Yes	212 (63.5)	20 (76.9)	1.9 (0.7–4.9)	0.154
Talwin/ritalin use				
No	228 (68.3)	16 (61.5)	1.0	
Yes	106 (31.7)	10 (38.5)	1.3 (0.6–3.1)	0.486
Morphine use				
No	209 (62.6)	21 (80.8)	1.0	
Yes	125 (37.4)	5 (19.2)	0.4 (0.1–1.1)	0.051
Heroin use				
No	299 (89.5)	26 (100)		
Yes	35 (10.5)	0 (0.00)	Not defined	
Injects daily				
No	52 (15.6)	5 (19.2)	1.0	
Yes	282 (84.4)	21 (80.8)	0.8 (0.3–2.1)	0.631
Injected at a hotel				
No	198 (59.3)	16 (61.5)	1.0	
Yes	136 (40.7)	10 (38.5)	0.9 (0.4–2.1)	0.821
injected at a shooting gallery				
No	287 (85.9)	17 (65.4)	1.0	
Yes	47 (14.1)	9 (34.6)	3.2 (1.4–7.7)	0.012
Ever used someone else's syringe				
No	136 (40.7)	2 (7.7)	1.0	
Yes	198 (59.3)	24 (92.3)	8.2 (1.9–35.5)	<0.001
Injected someone else as a service				

Table 3: Odds ratios and 95% confidence intervals for univariate analysis of HIV serostatus. (Continued)

No	256 (76.7)	13 (50.0)	1.0	
Yes	78 (23.3)	13 (50.0)	3.3 (1.5–7.4)	0.005
Obtains syringes from friends				
No	106 (31.7)	13 (50.0)	1.0	
Yes	228 (68.3)	13 (50.0)	0.5 (0.2–1.0)	0.063
Obtains syringes from questionable sources				
No	257 (77.0)	19 (73.1)	1.0	
Yes	77 (23.1)	7 (26.9)	1.2 (0.5–3.0)	0.658
Opposite-sex regular partners				
No	105 (31.4)	8 (30.8)	1.0	
Yes	229 (68.6)	18 (69.2)	1.0 (0.4–2.4)	0.944
Opposite-sex casual partners				
No	208 (62.3)	18 (69.2)	1.0	
Yes	126 (37.7)	8 (30.8)	0.7 (0.3–1.7)	0.474
Opposite-sex client partners				
No	290 (86.8)	19 (73.1)	1.0	
Yes	44 (37.7)	7 (26.9)	2.4 (1.0–6.1)	0.076
Regular same-sex partners				
No	326 (97.6)	24 (92.3)	1.0	
Yes	8 (2.4)	2 (7.7)	3.4 (0.7–16.9)	0.183
Casual same-sex partners				
No	320 (95.8)	25 (96.2)	1.0	
Yes	14 (4.2)	1 (3.9)	0.9 (0.1–7.2)	0.932
Client same-sex partners				
No	329 (98.5)	23 (88.5)	1.0	
Yes	5 (1.5)	3 (11.5)	8.6 (1.9–38.2)	0.013
Egocentric network variables (maximum number of network members = 20)				
Total number of IDU in network				
Median (IQR)	3 (2,5)	2 (0,5)	0.9 (0.8–1.1)	0.241
Total number of IDU family members in network				
Median (IQR)	0 (0,1)	0 (0,2)	1.2 (0.9–1.6)	0.260
Spouse IDU in network				
No	215 (64.4)	17 (65.4)	1.0	
Yes	119 (35.6)	9 (34.6)	1.0 (0.4–2.2)	0.917
IDU risk network variables (maximum number of IDU = 5)				
Number of IDU respondent sees daily				
None	107 (32.0)	12 (46.2)	1.0	
1	112 (33.5)	6 (23.1)	0.5 (0.2–1.3)	0.154
2	58 (17.4)	5 (19.2)	0.8 (0.3–2.3)	0.637
3–5	57 (17.1)	3 (11.5)	0.5 (0.1–1.7)	0.256
Number of IDU in network who inject daily				
None	191 (57.2)	17 (65.4)	1.0	
1	63 (18.9)	2 (7.7)	0.4 (0.1–1.6)	0.176
2	45 (13.5)	4 (15.4)	1.0 (0.3–3.1)	0.998
3–5	35 (10.5)	3 (11.5)	1.0 (0.3–3.5)	0.954
Number of IDU in network who have been IDU for >5 years				
0	159 (47.6)	14 (53.9)	1.0	
1	65 (19.5)	3 (11.5)	0.5 (0.1–1.9)	0.323

Table 3: Odds ratios and 95% confidence intervals for univariate analysis of HIV serostatus. (Continued)

2	52 (15.6)	4 (15.4)	0.9 (0.3–2.8)	0.819
3–5	58 (17.4)	5 (19.2)	1.0 (0.3–2.8)	0.969
Number of IDU in network who have injected at a hotel				
0	207 (62.0)	20 (76.9)	1.0	
1	64 (19.2)	2 (7.7)	0.3 (0.1–1.4)	0.135
2–5	63 (18.9)	4 (15.4)	0.7 (0.2–2.0)	0.458
Number of IDU in network who have injected at a shooting gallery				
0	254 (76.1)	20 (76.9)	1.0	
1	43 (12.9)	2 (7.7)	0.6 (0.1–2.6)	0.488
2–5	37 (11.1)	4 (15.4)	1.4 (0.4–4.2)	0.582
Number of IDU in network with whom respondent has pooled money				
0	94 (28.1)	12 (46.2)	1.0	
1	91 (27.3)	4 (15.4)	0.3 (0.1–1.1)	0.074
2–3	90 (27.0)	6 (23.1)	0.5 (0.2–1.5)	0.213
4–5	59 (17.7)	4 (15.4)	0.5 (0.2–1.7)	0.292
Number of IDU in network who have used a syringe before the respondent used it				
0	255 (76.4)	20 (76.9)	1.0	
1	56 (16.8)	4 (15.4)	0.9 (0.3–2.8)	0.869
2–5	23 (6.9)	2 (7.7)	1.1 (0.2–5.0)	0.894
Number of IDU in network who have used other drug paraphernalia before respondent has used it.				
0	201 (60.2)	17 (65.4)	1.0	
1	59 (17.7)	3 (11.5)	0.6 (0.2–2.1)	0.429
2–5	74 (22.2)	6 (23.1)	1.0 (0.4–2.5)	0.932
Number of IDU in network who have initiated respondent to injection drug use				
0	242 (72.5)	21 (80.8)	1.0	
1	77 (23.1)	5 (19.2)	0.7 (0.3–2.1)	0.563
2–5	15 (4.5)	0 (0.0)	Not defined	
Number of IDU in network who have shown respondent how to inject drugs				
0	229 (68.6)	19 (73.1)	1.0	
1	75 (22.5)	6 (23.1)	1.0 (0.4–2.5)	0.940
2–5	30 (9.0)	1 (3.9)	0.4 (0.1–3.1)	0.382
Number of IDU in network who have injected respondent with drugs				
0	161 (48.2)	12 (46.2)	1.0	
1	95 (28.4)	5 (19.2)	0.7 (0.2–2.1)	0.525
2–5	78 (23.4)	9 (34.6)	1.5 (0.6–3.8)	0.344
Number of IDU in network who have used talwin/ritalin				
0	239 (71.6)	17 (65.4)	1.0	
1–3	54 (16.2)	4 (15.4)	1.0 (0.3–3.2)	0.944
4–5	41 (12.3)	5 (19.2)	1.7 (0.6–4.9)	0.315
Number of IDU in network who have used cocaine				
0	210 (62.9)	18 (69.2)	1.0	
1	55 (16.5)	3 (11.5)	0.6 (0.2–2.2)	0.481
2–3	44 (13.2)	4 (15.4)	1.1 (0.3–3.3)	0.919
4–5	25 (7.5)	1 (3.9)	0.5 (0.1–3.6)	0.467
Number of IDU in network who have used morphine				
0	255 (76.4)	21 (80.8)	1.0	
1	46 (13.8)	4 (15.4)	1.1 (0.3–3.2)	0.924
2–5	33 (9.9)	1 (3.9)	0.4 (0.0–2.8)	0.336

Table 4: HCV multivariate adjusted OR and 95% CI.

Variable name	OR (95% CI)	p value
Years of injection drug use		
Categories based on quartiles (Q1–Q4 below)		
Q2 vs. Q1	3.1 (1.5–6.2)	0.002
Q3 vs. Q1	10.8 (4.8–24.5)	<0.001
Q4 vs. Q1	10.3 (4.7–22.6)	<0.001
injected at a shooting gallery		
Yes vs. no	4.3 (1.8–10.5)	<0.001
Ever used someone else's used syringe		
	See table 5	
Number of IDU in network who use talwin/ritalin		
	See table 5	
Number of IDU in network who have shown respondent how to inject drugs		
	See table 5	

After adjusting for these interactions, two respondent variables, obtaining needles from friends and injecting someone as a service, were no longer significant.

HBV was positively associated with three respondent variables: years of ID use, injection with a used syringe, and injection at a shooting gallery and two IDU risk network variables: the number of IDU in the risk network who injected talwin/ritalin, and the number who had injected at a hotel (table 6). The latter variable was significant only if the respondent identified 2 or more IDU in the risk network with this characteristic. HBV was negatively associated with reporting opposite-sex casual sex partners; moving to Winnipeg in the past year and the number of

IDU in the risk network who had initiated the respondent to injection drug use. As above, this variable was only significant if 2 or more IDU in the risk network were reported as having initiated the respondent to injection drug use.

HIV was positively associated with four respondent variables; injection with used syringes, injection at a shooting gallery, same-sex client sex partners and aboriginal ethnicity (table 7).

Discussion

Our two primary goals in this investigation were to determine which respondent and/or network variables were

Table 5: Adjusted odds ratios and 95% CI for interactions between injection with a used syringe and the IDU risk network variables.

A)				
		Reported injection with used syringe		
		Yes	No	
Number of talwin/ritalin users in IDU risk network	0	7.9 (3.7–16.8) 95 HCV +/58 HCV - (62%)	1.00 24 HCV +/81 HCV - (22.9%)	
	1–3	0.6 (0.3–42.4) 26 HCV +/10 HCV - (72%)	1.00 16 HCV +/7 HCV - (69.6%)	
	4–5	1.0 (0.2–156.8) 28 HCV +/9 HCV - (76%)	1.00 9 HCV +/2 HCV - (81.8%)	
	B)			
	Reported injection with used syringe			
			Yes	No
Number of IDU who have shown the respondent how to inject drugs	0	7.9 (3.7–16.8) 111 HCV +/41 HCV - (73.0%)	1.00 36 HCV +/63 HCV - (36.4%)	
	1	7.7 (0.3–37.1) 31 HCV +/23 HCV - (57.4%)	1.00 8 HCV +/20 HCV - (28.6%)	
	2–5	0.4 (0.2–87.2) 7 HCV +/13 HCV - (33.3%)	1.00 5 HCV +/7 HCV - (41.6%)	

(A) number of IDU in the network who used talwin/ritalin or (B) have shown the respondent how to inject. Serostatus data for HCV corresponding to each cell is also shown.

Table 6: HBV multivariate adjusted OR

Variable name	OR (95% CI)	p value
Years of injection drug use		
Categories based on quartiles (Q1–Q4 below)		
Q2 vs. Q1	4.7 (1.6–13.9)	0.005
Q3 vs. Q1	11.3 (3.8–33.2)	<0.001
Q4 vs. Q1	15.7 (5.3–46.5)	<0.001
Ever used someone else's used syringe		
Yes vs. No	2.2 (1.2–4.0)	0.005
injected at a shooting gallery		
Yes vs. no	2.9 (1.3–6.3)	0.020
Opposite-sex casual sex partners		
Yes vs No	0.5 (0.3–0.9)	0.020
Mobility (Moved to Winnipeg in past year)		
Yes vs. no	0.3 (0.2–0.7)	0.004
Number of IDU in network who use talwin/ritalin		
1–3 vs. none	2.9 (1.4–5.9)	<0.005
4–5 vs none	3.2 (1.4–7.2)	<0.009
Number of IDU in network who have initiated respondent to injection drug use		
1 vs. none	0.6 (0.3–1.4)	0.248
2–5 vs. none	0.1 (0.02–0.8)	0.023
Number of IDU in risk network who have injected at a hotel		
1 vs. none	1.3 (0.6–2.6)	0.490
2–5 vs. none	2.8 (1.3–5.9)	0.006

associated with HCV, HBV, and HIV in our setting and to compare and contrast these variables for the different pathogens. Only two respondent variables were positively associated with all three pathogens: injection with syringes previously used by another IDU and injection at a shooting gallery. The former represents a direct route for parenteral transmission of blood borne pathogens and is clearly a common factor for transmission of all three of the pathogens examined in our study area. The latter, given the high-risk activities characteristic of this type of venue [6,18,19], would be associated with an increased likelihood of exposure through purposeful or accidental use of contaminated equipment. Although this variable is a marker for high-risk behaviour on the part of individuals, it is also a proxy marker for membership in high risk, largely anonymous networks of IDU. Networks form around specific places where geographic place is the common connecting tie between individuals [22-24]. Including place in network analysis adds explanatory power for

understanding disease transmission regardless of the infection. Here shooting galleries are not simply high risk environments for individual users, but, by acting as repositories for contaminated equipment, they create bridging opportunities for pathogens between individuals who may not know each other and who may never otherwise meet.

Like shooting galleries, certain hotels in Winnipeg are also important venues for injection and favor formation of place-based networks of IDU. The likelihood of HBV infection increased proportional to the number of IDU the respondent knew who had injected at a hotel. Association with HBV, but not with HCV or HIV, could result from HBV entering the hotel network(s) earlier than HCV and HIV, resulting in more extensive spread. Alternatively, there may be specific behaviours associated with this group of IDU that favour specific transmission of this pathogen (e.g. hotel rooms, as opposed to shooting galleries, may offer more privacy and a greater likelihood of engaging in sexual activities, thus favouring HBV transmission over HCV). Regardless of the causal factors involved, like others [8,25], we found that incorporating interactions with risk network members provided a clearer understanding of what actually constitutes high-risk behaviour.

Respondents' injection of talwin/ritalin and their use by IDU network members was a marker for HCV and HBV. The type of drug used can loosely demarcate distinct networks of IDU [15,16]. Within these networks specific

Table 7: HIV multivariate adjusted OR

Variable name	OR (95% CI)	p value
Ethnicity		
Aboriginal vs. caucasian	4.4 (1.2–15.6)	0.022
Ever used someone else's used syringe		
Yes vs. No	8.7 (2.0–38.5)	0.004
injected at a shooting gallery		
Yes vs. no	3.0 (1.2–7.5)	0.018
Same-sex client sex partners		
Yes vs No	6.9 (1.5–32.4)	0.014

behaviours may favour transmission (e.g. the room temperature production of cocaine *vs.* the need to heat some other types of drugs [14]). To our knowledge, extensive use of talwin/ritalin is relatively rare outside of the Canadian provinces of Manitoba and Saskatchewan. It is notable that in our analysis, injection with used syringes interacted with talwin/ritalin such that the prevalence of HCV was high amongst talwin/ritalin users regardless of whether or not they reported injection with used syringes. Like cocaine, talwin/ritalin is typically prepared at room temperature and frequently involves communal use of the prepared drug and filters, which may account for the high prevalence of HCV regardless of syringe sharing practices (typically pills are crushed in room temperature water; IDU then use a common filter to draw drug into their own respective syringes). It is also notable, that HIV does not show a positive association with talwin/ritalin in contrast to HCV and HBV. As above, this may reflect a different temporal framework for pathogen entry to the network of talwin/ritalin users. If this is the case, the same behaviours which facilitated spread of HBV and HCV may ultimately lead to high HIV prevalence in this group.

Two IDU risk network variables were negatively associated with either HCV or HBV. For HBV, there was a negative association with the number of IDU in the respondent's risk network who had initiated them to injection drug use, while for HCV there was a negative association if the respondent reported that there were IDU in their risk network who had shown them how to inject drugs. For HCV, there was also a weak interaction between the number of IDU members who had shown the respondent how to shoot drugs and syringe sharing, such that syringe sharing was no longer positively associated with prevalence. In general, these network variables may mark small, relatively closed networks of IDU who may share equipment and syringes, but rarely admit new members. Low network turnover has been associated with lower risk [12] as opportunities for entry of pathogens to these networks (via infected individuals) would be correspondingly reduced. An area worthy of research would be whether the lower prevalence of infection is a chance byproduct of the social interactions within these networks or whether the individuals within closed networks are actively and knowingly attempting to isolate themselves from infection risk.

The greater likelihood of HIV and HBV being transmitted through sexual contact was consistent with sex partner-related variables being retained in the final models for these two pathogens. However, while a positive correlation for HIV occurred with same-sex client partners, the directionality for HBV and opposite-sex casual partners was opposite. Although the direction of this relationship appears counter-intuitive, it may reflect a greater likeli-

hood to engage in safer sex harm reduction activities with sex partners of this type. This question could be addressed in more detailed research on sexual contact behaviours in this population.

Aboriginal ethnicity and a recent move to Winnipeg were associated with HIV and HBV, respectively. HIV was the only pathogen where ethnicity remained as an independent predictor of risk in the final model. Research in other cities have identified specific social network interactions that can account for ethnic differences in disease prevalence [13]. Further research would be necessary within our study population to identify what risk behaviour(s) may be associated with HIV infection in the aboriginal population in Manitoba that may not have been captured by our survey. The relatively small number of HIV positive individuals in our dataset also may have limited the analytic potential for this pathogen in regression analysis. A recent move to Winnipeg was associated with a lesser likelihood of HBV positivity. This pattern may reflect a lower prevalence of HBV and/or earlier or more extensive deployment of vaccination programs in the provinces from which these IDU originated (e.g. the province of British Columbia began vaccinating approximately 6 years prior to Manitoba).

A number of limitations for the study should be noted. First, the study was cross-sectional and hence does not provide any data on trends or whether currently observed behaviours were a result of, rather than the cause of, an infection by one of the pathogens studied. Second, the data were egocentric and hence no independent validation of the behaviours of IDU in the risk network was possible. Third, network members were elicited based on contact in the previous 30 days, but questions regarding individual behaviours and interactions with network members were typically collected for the previous 6 months (or longer periods for some variables such as "ever-use of someone else's syringe"). Time frames are relevant, given the noted lack of correlation between some known risk behaviours and serostatus in our analysis (see paragraph below). Any comparisons of our results with those from other areas would need to take note of the time frames used. Fourth, the egocentric and IDU risk networks were truncated at 20 and 5 people, respectively. The implications of this data truncation are greatest for the IDU risk network variables as they formed a large part of the analysis. Overall, 17% of respondents reported more than 5 IDU in their egocentric network. We felt that full data collection on all IDU in the egocentric network was impractical in terms of time, complexity of data collection and accurate recall by respondents, however, it should be noted that this additional data could have revealed additional patterns not apparent in our results.

Finally, some limitations regarding specific variables should be noted. First, data on sharing of drug-preparation equipment and transfer of prepared drugs between syringes was available for only the six months prior to interview. In contrast, sharing of used syringes was based both on six-month and "ever" data. Only the latter was a predictor of risk, therefore, if more extensive data had been collected for equipment sharing and drug transfers, they may also have been significant predictors of risk of infection for some, or all, of the pathogens studied. Second, only a limited number of sexual behaviour questions were included in the study and no specific questions on type of sex were asked. Collection of data on anal sex, in particular, could have revealed other correlations not evident with the dataset available. Third, only "injecting someone else as a service" was analyzed; "injecting someone else as a favor" or "receiving injections from others" can be considered distinct traits and could also be related to the common use of syringes and pathogen transmission.

Conclusion

Our findings reinforce the clear link between respondent behaviours (e.g. injection with someone else's used syringe) and risk of infection. However, they also highlight the importance of defining the types of personal networks that form within an area and identifying the different patterns of transmission of pathogens within those networks. In our setting, pathogen transmission is associated with loosely defined, sometimes anonymous, high-risk networks formed around specific venues (shooting galleries, hotels) or between users who are linked by their drug use preferences. Smaller, somewhat isolated pockets of IDUs appear to exist within the larger population where behavioural patterns pose a lesser risk, unless, or until, a given pathogen enters those networks. Temporal differences with respect to pathogen introduction to the different networks likely influences which specific behaviours appear as protective or as high-risk, highlighting the importance of verifying the extent to which risk behaviours or markers identified as important in one setting or to one pathogen apply to other settings or pathogens. The lack of any apparent correlation between the prevalent infections for the three pathogens, noted in the results section, reinforces the need to identify the different social and temporal patterns associated with the various pathogens that may be circulating within a population. Both individual- and network-level concepts assist in characterizing and comparing the transmission of HCV and HBV, as well as, HIV.

A potential hierarchy of risk emerges, defined by space and time. Individuals may have some independence in their decisions to use drugs or not, or share syringes or not. However, the broader personal network in which

they find themselves, whether defined by living near a hotel where drug use is common or by their ethnic or age group, is no less important in contributing to their potential for initiation of specific behaviours or exposure to infectious agents. The smaller, more specific social interactions with their IDU risk network forms an additional step in a progression which may determine an individual's choice of drug; their norms, rituals and/or practices of injection; and their likelihood of exposure to infected individuals. Both individual- and network-level concepts assist in characterizing and comparing the transmission of HCV, HBV, and HIV and determining the underlying patterns that drive the social connections between individuals that may favour or hinder transmission of specific pathogens.

Competing interests

The author(s) declare that they have no competing interests.

Authors' contributions

JW and AJ conceived and designed the study. JW oversaw implementation of the study in Winnipeg and drafted the manuscript. LS prepared the database for analysis and both JW and LS performed the statistical analysis. AJ and LS assisted with manuscript editing and all authors read and approved the final manuscript.

Acknowledgements

The authors wish to thank Debbie Nowicki for assisting in implementation of the study, Dr. Magdy Dawood for facilitating specimen testing, Dr. Sam Friedman for advice on questionnaire development and Souradet Shaw for advice on analysis. The study was funded by an operating grant from the Canadian Institutes of Health Research.

References

1. Klovdahl AS: **Social networks and the spread of infectious diseases: the AIDS example.** *Soc Sci Med* 1985, **21**:1203-1216.
2. Klovdahl AS, Potterat JJ, Woodhouse DE: **Social networks and infectious disease: the Colorado Springs study.** *Soc Sci Med* 1994, **38**:79-88.
3. Fairbairn N, Wood E, Small W, Stoltz J, Li K, Kerr T: **Risk profile of individuals who provide assistance with illicit drug injections.** *Drug Alcohol Depend* 2005, **82**:41-46.
4. Hahn JA, Page-Shafer K, Lum PJ, Ochoa K, Moss AR: **Hepatitis C virus infection and needle exchange use among young injection drug users in San Francisco.** *Hepatology* 2001, **34**:180-187.
5. Friedman S, Furst RT, Jose B, Curtis R, Neaigus A, Des Jarlais DC, Goldstein M, Ildefonso G: **Drug scene roles and HIV risk.** *Addiction* 1998, **93**:1403-1416.
6. Friedman S, Kang SY, Robles RR, Colon HM, Andia JF, Oliver-Velez D, Finlinson A: **Drug-scene roles and HIV risk among Puerto Rican injection drug users in East Harlem, New York and Bayamon, Puerto Rico.** *J Psychoactive Drugs* 2002, **34**:363-369.
7. Latkin CA, Forman V, Knowlton A, Sherman S: **Norms, social networks, and HIV-related risk behaviours among urban disadvantaged drug users.** *Soc Sci Med* 2003, **56**:465-476.
8. Neaigus A, Friedman S, Jose B, Goldstein M, Curtis R, Ildefonso G, Des Jarlais DC: **High-risk personal networks and syringe sharing as risk factors for HIV infection among new drug injectors.** *J Acquir Immune Defic Syndr Hum Retrovirology* 1996, **11**:499-509.
9. Needle RH, Coyle S, Cesari H, Trotter R, Clatts M, Koester S, Price L, McLellan E, Finlinson A, Bluthenthal MA, Pierce T, Johnson J, Jones TS, Williams M: **HIV risk behaviours associated with the injection**

- tion process: multiperson use of drug injection equipment and paraphernalia in injection drug user networks.** *Subst Use Misuse* 1998, **33**:2403-2423.
10. Latkin CA, Mandell W, Vlahov D, Oziemkowska M, Celentano D: **People and places: behavioural settings and personal network characteristics as correlates of needle sharing.** *J Acquir Immune Defic Syndr Hum Retroviro* 1996, **13**:273-280.
 11. Klee H, Morris J: **The role of needle exchanges in modifying sharing behaviour: cross-study comparison 1989-1993.** *Addiction* 1995, **90**:1635-1645.
 12. Hoffman JP, S. SS, Pach A: **Changes in network characteristics and HIV risk behavior among injection drug users.** *Drug Alcohol Depend* 1997, **46**:41-51.
 13. Kottiri BJ, Friedman S, Neaigus A, Curtis R, Des Jarlais DC: **Risk networks and racial/ethnic differences in the prevalence of HIV infection among injection drug users.** *J Acquir Immune Defic Syndr Hum Retroviro* 2002, **30**:95-104.
 14. Grund JC, Friedman S, Stern LS, Jose B, Neaigus A, Curtis R, Des Jarlais DC: **Syringe-mediated drug sharing among injecting drug users: patterns, social context and implications for transmission of blood-borne pathogens.** *Soc Sci Med* 1996, **42**:691-703.
 15. Crofts N, Hopper JL, Milner R, Breschkin AM, Bowden DS, Locarnini SA: **Blood-borne virus infections among Australian injecting drug users: implications for spread of HIV.** *Eur J Epidemiol* 1994, **10**:687-694.
 16. Krekulova L, Rehak V, Madrigal N, Johnson M, Killoran P, Riley LW: **Genotypic and epidemiologic characteristics of hepatitis C virus infections among recent injection drug user and non-user populations.** *Clin Infect Dis* 2001, **33**:1435-1438.
 17. Paschane DM, Fisher DG: **Etiology of limited transmission diseases among drug users: does recent migration magnify the risk of sharing injection equipment?** *Soc Sci Med* 2000, **50**:1091-1097.
 18. Reyes JC, Robles RR, Colon HM, Freeman DH, Sahai H, Matos TD: **Risk factors for shooting gallery use among drug injectors in Puerto Rico.** *P R Health Sci J* 1996, **15**:227-231.
 19. Santibanez SS, Garfein RS, Swartzendruber A, Kerndt PR, Morse E, Ompad DC, Strathdee S, Williams IT, Friedman SR, Ouellet LJ: **Prevalence and correlates of crack-cocaine injection among young injection drug users in the United States.** *Drug Alcohol Depend* 2005, **77**:227-233.
 20. Hosmer DW, Lemeshow S: **Applied Logistic Regression.** 2nd edition. New York, John Wiley & Sons, Inc.; 2000:375.
 21. Tabachnick BG, Fidell LS: **Using multivariate statistics.** Needham Heights, MA, Allyn & Bacon; 2001:966.
 22. Klodahl AS, Graviss EA, Yaganehdoost A, Ross MW, Wanger A, Adams GJ, Musser JM: **Networks and tuberculosis: an undetected community outbreak involving public places.** *Soc Sci Med* 2001, **52**:681-694.
 23. Potterat JJ, Rothenberg R, Woodhouse DE, Muth SQ, Pratts CI, Fogle JS: **Gonorrhoea as a social disease.** *Sex Transm Dis* 1985, **12**:25-32.
 24. De P, Singh AE, Wong T, Yacoub W, Jolly AM: **Sexual network analysis of a gonorrhoea outbreak.** *Sex Transm Infect* 2004, **80**:280-285.
 25. Friedman SR, Neaigus A, Jose B, Curtis R, Goldstein M, Ildefonso G, Rothenberg R, Des Jarlais DC: **Sociometric risk networks and risk for HIV infection.** *Am J Public Health* 1997, **87**:1289-1296.

Pre-publication history

The pre-publication history for this paper can be accessed here:

<http://www.biomedcentral.com/1471-2458/6/229/prepub>

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

